

Environmental Information Sheet

FOLICUR® MAPP number 11278



An emulsion (oil in water) formulation containing 250 g/l (25.9% w/w) tebuconazole
 A broad spectrum systemic fungicide for winter and spring crops of wheat (excluding durum), barley, oats, rye and oilseed rape; also field beans, cabbages, carrots, horseradish, leeks, linseed, parsnips, swedes and turnips.

Maximum individual dose: 1.0 litres product per hectare for all crops

Maximum total dose: 1 litre/ha for linseed,
 2 litre/ ha for cereals, field beans, swedes and turnips,
 2.25 litres/ha for cabbages,
 2.5 litres/ha for oilseed rape,
 3 litres/ha for carrots, parsnips, horseradish and leeks

Section	Profile
<p>1. WILDLIFE</p> <p>Mammals and Birds</p>	<p>Folicur® Is not classified as '<i>Harmful to game and wildlife</i>'</p> <p>No risk management is necessary to protect mammals and birds. Folicur® is of low/moderate toxicity to mammals and birds. The risk to wild mammals and birds grazing on treated areas is low, as is the risk due to exposure from other routes, e.g. consumption of insects.</p>
<p>2. BEES</p>	<p>No risk management necessary.</p> <p>Folicur® is of low risk to honeybees</p>
<p>3. NON TARGET INSECTS AND OTHER ARTHROPODS</p>	<p>No risk management necessary.</p> <p>Folicur® poses a low risk to a range of arthropod species commonly found in and around treated fields, e.g. ground beetles, lady birds and aphid parasitoids.</p>
<p>4. AQUATIC LIFE</p>	<p>Folicur® is classified as "<i>HARMFUL TO FISH OR OTHER AQUATIC LIFE</i>"</p> <p>Tebuconazole is "<i>TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.</i></p> <p>Folicur® is moderately toxic to fish, aquatic invertebrates and algae. However, no risk management is necessary. When used according to the label instructions there will not be sufficient contamination of water to present a risk to aquatic life:</p> <p><i>DO NOT CONTAMINATE SURFACE WATERS OR DITCHES WITH CHEMICAL OR USED CONTAINER.</i></p> <p>Folicur® is not classified under the LERAP scheme</p>

Environmental Information Sheet

FOLICUR® MAPP number 11278



Section	Profile
5. SOIL and GROUNDWATER	Tebuconazole is moderately persistent in soil, but has low mobility in soil. Use of Folicur® according to the label presents a low risk to groundwater.
Earthworms	Tebuconazole is of low toxicity to earthworms. No risk management is necessary, when Folicur® is used according to the label.
Soil Micro-organisms	Folicur® is of low risk to soil micro-organisms. No risk management necessary in order to prevent risks to the processes of soil respiration and nitrogen turnover.
6. NON-TARGET PLANTS	When used as recommended Folicur® is not expected to have adverse effects on non-target plants.

ALWAYS READ THE LABEL: USE PESTICIDES SAFELY

For further information about the environmental profile of this product contact:-

Bayer CropScience Limited
Hauxton
Cambridge
CB2 5HU

Bayer CropScience Assist Technical Help-Line 0845 6092266
This Environmental Information sheet was prepared in accordance with CPA Guidance Notes Version 3

e-mail: tech@bayercropscience.com

© EIS, copyright of Bayer CropScience Limited